

## A CONCEPTUAL FRAMEWORK FOR ASSESSING THE FIELD EFFICIENCY OF DRONES IN IDENTIFYING POTENTIAL BREEDING SITES OF THE *Aedes* MOSQUITO

Zulfadli Mahfodz<sup>1,2\*</sup>, Nazri Che Dom<sup>1</sup>, Hasber Salim<sup>3</sup> and Nopadol Precha<sup>4</sup>

<sup>1</sup>Faculty of Health Sciences, Universiti Teknologi MARA, Selangor Branch, Puncak Alam Campus, 42300 Selangor, Malaysia

<sup>2</sup>Faculty of Applied Sciences, Universiti Teknologi MARA, Perak Branch, Tapah Campus, Tapah Road, 35400 Perak, Malaysia

<sup>3</sup>School of Biological Sciences, Universiti Sains Malaysia, 11800 Penang, Malaysia

<sup>4</sup>Department of Environmental Health and Technology, School of Public Health, Walailak University, Nakhon Si Thammarat, Thailand

### ARTICLE INFO

#### Keywords:

*Aedes mosquito,*  
*breeding site,*  
*drones,*  
*machine learning,*  
*surveillance*

### ABSTRACT

*The identification of breeding sites is crucial for effective dengue prevention strategies. Community readiness and surveillance of these sites are essential for controlling the Aedes mosquito population. Drones have emerged as a promising tool for surveillance activities. This paper aims to develop a conceptual framework and present comprehensive intervention methods for controlling dengue cases. The study explores community readiness, habitat profiling, and mapping. Phase I focuses on community readiness, using a questionnaire to gather information about the community's knowledge, attitudes, and practices related to dengue control. Phase II involves profiling and mapping potential breeding sites of Aedes mosquitoes in selected areas using innovative drone technologies. This phase aims to identify and characterize breeding sites based on various parameters, such as size, water quality, and proximity to human dwellings. Systematic aerial surveys with drones equipped with high-resolution cameras and sensors will capture detailed images and environmental data. The results will be presented in interactive maps and detailed reports, supporting targeted interventions and data-driven decision-making for effective dengue prevention and control. This conceptual framework can assess the efficiency of drones as an alternative tool for dengue surveillance, which can be applied in dengue-endemic regions.*

## 1. INTRODUCTION

*Aedes aegypti* is the main vector of dengue, yellow fever, and chikungunya (Passos et al., 2022). There are no vaccines or antiviral treatments available to treat mosquito-borne diseases. Therefore, the most appropriate method currently available for combating these diseases is to remove any potential mosquito breeding sites (Rahman et al., 2021). In general, the breeding sites can be categorized as either artificial or natural containers (Amarasinghe et al., 2017). *Aedes* mosquitoes reproduce in stagnant and clean water (Mehra et al., 2016). Subsequently, all water-holding containers have the potential to serve as breeding grounds. Mosquito control and monitoring efforts can become costly, time-wasting, and ineffective without sufficient technical support (Passos et al., 2022). Contributing to the failure of control programmes are human social factors, improper garbage collection, and a lack of hygienic

awareness (Hasnan et al., 2017; Madzlan et al., 2017). The strategies employed to control mosquitoes differ according to the individual, community, and regional environments (Joshi & Miller, 2021). The COVID-19 pandemic has affected dengue monitoring and control field technicians dynamically as they confront a new occupational hazard (Valdez-Delgado et al., 2021). Currently, health inspectors are required to visit residences to identify and destroy mosquito breeding grounds. This practice is difficult for officials to carry out and presents a number of disadvantages, including time limits, safety hazards, and high expenses (Mehra et al., 2016; Passos et al., 2022). Drones were originally envisioned as simple devices, but their complexity has developed in tandem with the intricacy of their designated duties. According to Hardy et al. (2022) and Mohd Daud et al. (2022), the type, size, power, and application conditions of drones are the main factors that determine their operational capabilities.

\*Corresponding author: zulfa2015@uitm.edu.my

Drones are used in public health research for a wide range of purposes, including tracing survivors after natural disasters, shipping medical supplies to isolated areas, and providing initial treatment in emergency locations (Carrillo-Larco et al., 2018; Fornace et al., 2014; Hiebert et al., 2020; Passos et al., 2022). Drones are also put into operation in various other fields, including agriculture, forestry, ecology, and environmental monitoring (Stanton et al., 2021). The evidence on the efficiency of drones in finding mosquito breeding sites is insufficient and limited (Valdez-Delgado et al., 2021). Satellite imaging is considered unfeasible due to its long repeat times, cloud contamination, limited geographical and temporal resolutions, and high costs (Fornace et al., 2014; Hardy et al., 2022). Rapid technological advancements, encouraged by commercial demand and open market competition, have changed drones from mere hobbies into potentially useful tools that can shorten surveillance activities and enrich mosquito control programmes (Faraji et al., 2021).

In this study, aerial photography of water retention from drones used to produce a map. It supplies data based on the shadow effect and the tilt angle of the drone camera with adequate precision to detect water retention. This is related to the studies in Carrasco-Escobar et al. (2019) and Haas-Stapleton et al. (2019), which examine the properties of water bodies to identify potential mosquito breeding places. In the study conducted by Dias et al. (2018), drones were operated to attain many aerial image configurations for a database. Using the collected photos and the annotated database, the proposed system was tested and trained. This technology allows the detection of tiny objects that cannot be observed by existing remote sensing methods (Bravo et al., 2021). In the study, Mukabana et al. (2022) provided an example of architectural innovation by combining two established technologies and executing them in a new market and context. The results revealed that a combination of drone application and insecticides can effectively reduce mosquito populations in an irrigated rice agroecosystem. Most of the research emphasizes that developments in computer and software engineering and computer science are the primary drivers of drone technological growth (Mohd Daud et al., 2022). Numerous significant outcomes have been yielded by the application of technology to drones. Drone surveying attempts for man-made containers, for instance, are made more precise by pairing Global Positioning System (GPS) receivers with machine learning techniques and image technologies like multispectral imaging (Schenkel et al., 2020). In the following investigation was reported in Ali et al. (2022), they merged the Internet of Medical Things (IoMT) and Geographic Information System (GIS) maps. The authors provide a way to mitigate and manage dengue virus outbreaks through call data record analysis. Once the patient's specific location has been confirmed, a spray unit will be alerted to send out drones to treat the affected area. In another scenario, Hardy et al. (2022) analyse drone data using a hybrid of mapping methods: supervised image classification using machine learning and technology-assisted digitising mapping that is accessible even

to those with no technical background. As stated in Valdez-Delgado et al. (2021), the advancement of machine learning must be further investigated, and further research is required to spread out the usage of neural networks in mosquito surveillance. Improving the precision of technology requires a comprehensive commitment and data from many images representing a vast array of situations.

As preliminary data, we conducted a SWOT analysis (Figure 1) on the viability of using drones for surveillance based on information from related articles. Each article described a new and better way to use drone images to find possible mosquito breeding grounds. The ability of drones to fly in difficult-to-access areas is their greatest advantage. According to Hardy et al. (2022), drones can not only capture images or videos with greater spatial or temporal resolution but also from a variety of angles and heights. In addition, deploying programmable drones increases health inspectors' safety by decreasing their contact with potentially hazardous incidents or sets. The initial cost of drone hardware is considerable, but it is feasible since a small team can cover a vast area from the air with minimal operating costs (Chiroli et al., 2017). Drone implementation for mosquito breeding site surveillance may face numerous challenges. For example, the laws and guidelines for flying drones in each region are different. There is no standard set of guidelines for operating or regulating drones; there are restricted zones where drones cannot be used; there are topographic and climate differences; there are privacy concerns; and community attitudes towards and acceptance of drones vary across cultures (Annan et al., 2022; Mohd Daud et al., 2022; Poljak & Šterbenc, 2020). The scalability of the proposed method may be limited due to limitations in the geographical area that a single drone can cover (Bravo et al., 2021). However, Hardy et al. (2017) demonstrated that a low-cost drone can be used to map water bodies in various settings, including natural water bodies, rice paddies, and urban areas, with the goal of identifying and mapping aquatic mosquito habitats. A well-organized database for identifying and categorizing objects in aerial videos is also an essential element. Among the features highlighted by Passos et al. (2022), a database should have a significant number of samples for each target group, an absence of image distortions, and precise object annotation.

Detecting water bodies in remote areas using drone-captured images requires several technological and competence hurdles. Creating orthomosaics from drone imagery is a time-consuming process that calls for a powerful computer, a huge space for records of information, and specialised software (Stanton et al., 2021; Wyngaard et al., 2018). In addition to hardware and software concerns, misconfigured drones can pose risks and vulnerabilities (Wyngaard et al., 2018). Flight experience was essential for calculating the ideal flight hours, as the drone could not be piloted in rainy weather and the influence of adverse weather on the equipment could cause it to overheat on hot days. More engagements between stakeholders and specialists across disciplines are required to provide more specific recommendations on how this technology can be used most efficiently (Stanton et al., 2021).



**Figure 1:** SWOT analysis on the viability of using drones for surveillance activities

Therefore, from the literature search and SWOT analysis, the objective of this paper is to develop a conceptual framework and discuss the intervention methods that are summarized in Figure 2. These phases revolve around three main interventions whereby community assessment, profiling potential habitat, and technology integration are fully taken into consideration in developing sustainable dengue preventive measures. It is essential to have a comprehensive understanding of the conceptual frameworks associated with different aspects in dengue management. This includes understanding how communities are mobilized, their roles in vector control, efforts in health education, and the ability to sustain new prevention measures over a long period of time.

**2. PHASE I: COMMUNITY ASSESSMENT**

Many studies related to the prevention and control of the *Aedes* mosquito indicate that household intervention can aid in larval reduction programmes; however, empowering and sustaining such practises over time and outside of research projects can be challenging (Rahman et al., 2021)we aimed to map the spatial distribution of female adult *Ae. aegypti* and predict its abundance in northeastern Thailand based on socioeconomic, climate change, and dengue knowledge, attitude and practices (KAP. The integration of household involvement into routine dengue programmes, including the most effective strategies, types of educational materials needed, programme regularity, and participation models, is still the subject of ongoing research. As discussed by Kleinman (2010), interventions are dynamic, can result in unforeseen effects, and are socially established. A review of dengue programmes in four countries by Horstick et al. (2010)questionnaires indicated several factors that contributed to the failure of the programmes. It includes a shortage of manpower, funds, and knowledge; an overdependence on chemicals; a low level of household participation; and a nearly non-existent comprehensive assessment system. Therefore, any interventions must be tailored according to the local landscape and engage specific communities at risk (Carrasco-Escobar et al.,

2022)global health security has been threatened by the geographical expansion of vector-borne infectious diseases such as malaria, dengue, yellow fever, Zika and chikungunya. For a range of these vector-borne diseases, an increase in residual (exophagic.

**2.1 Evaluation of the Conceptual Framework**

This phase focuses on assessing community readiness for dengue prevention strategies using a detailed questionnaire. The questionnaire collects data on community demographics, knowledge of dengue (such as awareness of transmission and symptoms), attitudes towards dengue prevention (including concern and perceived responsibility), and practices related to dengue control (such as source reduction and use of protective measures). It also explores perceived barriers and the community’s acceptance of drone technology for mosquito surveillance. This data is coded and analyzed to identify common themes and develop a theoretical framework that explains community readiness and influencing factors. This approach ensures that interventions are culturally sensitive and contextually appropriate, enhancing community participation and supporting effective dengue prevention implementation. The insights gained help tailor interventions to specific community needs and foster a collaborative approach to dengue control (Charmaz, 2014; Creswell & Poth, 2018).

**2.2 Feasibility of the Approach**

To enhance the effectiveness and sustainability of dengue control measures, it is crucial to promote evidence-based strategies, utilize local knowledge and scientific skills, and incorporate innovative solutions in collaboration with communities. Implementing new technologies is most convenient when linked with participatory processes that strengthen household engagement, civil society, and system accountability. Participatory action research framework can be used to reevaluate any challenges and rapidly test multiple strategies tailored to specific geographic locations (Duque et al., 2019). To enable the effective implementation of health data systems, specialized techniques are specifically designed for healthcare domains. Thus, enhance the performance and effectiveness of the system, facilitate learning and user experience, minimize medical faults, design time, funding, and training costs (Zhang, 2005). This approach will also emphasize the performance efficiency, security and privacy, usability, and portability of the proposed project. This process will ensure that the technology products adopted for the project are prototyped, examined, and fine-tuned to improve their performance and acceptance by end-users.

**3. PHASE II: PROFILING POTENTIAL BREEDING SITES OF *Aedes* MOSQUITO**

In several regions worldwide, *Aedes* mosquitoes primarily inhabit artificial containers (Carrasco-Escobar et al., 2022; Madzlan et al., 2017). The physical landscape characteristics and availability of containers that can retain water have been discovered to be critical elements in the reproduction of mosquitoes in Malaysia and other countries (Muñiz-sánchez et al., 2022). Knowing the habitat

characteristics and other elements like vegetation indices, land use, canopy cover, and elevation can provide comprehensive knowledge about mosquito abundance and distribution (Carrasco-Escobar et al., 2022). Health inspectors are currently conducting larval surveys and destroying dengue breeding grounds in dengue hotspot areas as part of the Ministry of Health Malaysia's preventative measures. Due to broad coverage areas and a limited number of health inspectors in each district, this task is time-consuming and cannot be completed properly. Consequently, drone-based monitoring of breeding sites will have a wider scope, be more prospective, and require less human work.

### 3.1 Drone Surveillance

Aerial surveillance permits the identification of potential breeding sites in areas that are inaccessible to traditional ground surveillance. Thus, it assists health inspectors in accurately identifying and mapping the breeding sites using drone images (Muñiz-sánchez et al., 2022). The size of mosquito breeding sites can be smaller than 1 meter; therefore, the ability of drones to capture high-precision images is critical for analysis (Hardy et al., 2022). The drone surveillance will be conducted using the method explained in Carrasco-Escobar et al. (2019). In selected study areas, weekly aerial surveys will be conducted using a drone to identify suitable outdoor breeding places such as pails, vessels, bins, and bowls. Any man-made structures outside the buildings that can serve as breeding grounds directly or indirectly will also be inspected. The drone will fly to an altitude of approximately 100 meters in the selected urban and rural areas, resulting in a sample distance of about 0.1 meters per pixel on the ground. To construct an orthomosaic, 100 waypoints will be automatically determined in each grid to give a 70% overlap between neighbouring images.

### 3.2 Entomological Surveillance

Entomological surveillance is crucial to effectively monitor and assess interventions, identify regions that require further surveillance, and assess the possibility of vector-borne transmission (Carrasco-Escobar et al., 2022)global health security has been threatened by the geographical expansion of vector-borne infectious diseases such as malaria, dengue, yellow fever, Zika and chikungunya. For a range of these vector-borne diseases, an increase in residual (exophagic. The proposed method is modified from Muñiz-sánchez et al. (2022). The *Aedes* larval surveillance data will be derived from weekly inspections of selected areas for both natural and artificial containers. A breeding site will be georeferenced if a positive *Aedes* mosquito is identified or suspected. A breeding site will be considered positive if the sample contains at least one *Ae. aegypti* or *Ae. albopictus* larva. Expert taxonomists will identify the *Aedes* larvae by using morphological criteria. Due to the varying sizes and forms of the research area, this study will superimpose a 200-meter grid system on the map of the study area and use these fixed grid cells as study units to minimise the amount of normalisation required. Additionally, surveillance will be conducted within a 200-meter radius of the gravid oviposition traps. After five days of

placement, the traps will be returned to the laboratory. Concurrently, larval surveys will be done at prospective breeding sites identified by the drone survey. The larvae of mosquitoes will be identified to the species level using the criteria given by the Malaysian Ministry of Health. Three larval indices, namely the *Aedes* Index, Container Index, and Breteau Index, will also be measured in accordance with World Health Organization (WHO) standards.

### 3.3 Data Analysis and Image Classification

The images captured by the drone will be used for visual inspection to find any water bodies that may serve as potential breeding sites. Subsequently, a comprehensive map of the mosquito breeding grounds and the surrounding landscape can be generated (Carrasco-Escobar et al., 2022)global health security has been threatened by the geographical expansion of vector-borne infectious diseases such as malaria, dengue, yellow fever, Zika and chikungunya. For a range of these vector-borne diseases, an increase in residual (exophagic. Drones can generate various maps, including two-dimensional maps, elevation maps, thermal maps, and three-dimensional maps or models. It can also gather precise information about the landscape with a ground resolution that is significantly higher than that of any current satellite imaging (Budiharto et al., 2021)developing 3D models using photogrammetric and situation mapping uses geographic information systems. The drone used has advantages in a wider range of areas with adequate power support. The drone is also supported by a high-quality camera with dreadslocks for image stability, so it is suitable for use in mapping activities. Conclusions: Using Google earth data at two separate locations as a benchmark for the accuracy of measurement of the area at three variations of flying height in taking pictures, the results obtained were 98.53% (98.68%. AgiSoft PhotoScan Pro will be utilized for photogrammetric analysis, which requires photographically based surface measurements (Hardy et al., 2017). The generated drone photography will be imported into PhotoScan and used to construct orthomosaics, which are geo-referenced mosaics of overlapping images with topographic distortion correction for each study site. The photogrammetric method can be utilized to create a three-dimensional landscape that contains both natural and man-made parts of the environment. Then, it can be fed straight into Geographic Information System (GIS) software for visualisation (Carrasco-Escobar et al., 2022)global health security has been threatened by the geographical expansion of vector-borne infectious diseases such as malaria, dengue, yellow fever, Zika and chikungunya. For a range of these vector-borne diseases, an increase in residual (exophagic. The potential breeding sites will be categorized based on various parameters, such as size, water quality, and proximity to human dwellings.

### 3.4 Machine Learning

Integration of available technologies like drones, machine learning, deep learning and big data, can be used to have dependable evidence about mosquito population dynamics and administering areas at risk (Muñiz-sánchez et al., 2022). Machine learning demonstrates a crucial role in identifying mosquito breeding sites by utilizing

collected data to train and classify ecological and geographical landscape associated with potential breeding sites. Machine learning models, although are not perfect, can differentiate between breeding and non-breeding sites by processing large datasets using algorithms such as Random Forest and Support Vector Machines (SVM) (Joshi & Miller, 2021). Deep learning approaches like convolutional neural networks (CNNs) are effective in image recognition (Bravo et al., 2021). It is anticipated that the prediction models would reasonably recognise proper images of potential breeding sites with water retention zones. This approach allows local authorities to manage mosquito control efforts more effectively, thus reducing dengue cases.

Machine learning approaches have made a significant contribution, especially in mapping potential breeding sites for mosquitoes. In the study by Hardy et al. (2022), several dominant land cover classes were used, such as open water, open water with sunlight, emergent vegetation, dry, and land cover types, to assist in the classification process. In another example, Rahman et al. (2021) we aimed to map the spatial distribution of female adult *Ae. aegypti* and predict its abundance in northeastern Thailand based on socioeconomic, climate change, and dengue knowledge, attitude and practices (KAP used five supervised learning models, which are logistic regression, support vector machine, k-nearest neighbour, artificial neural networks, and random forest based on socioeconomic, climate change, and dengue knowledge, attitude, and practices (KAP), to predict the abundance of mosquitoes (high and low). The use of machine learning models for the prediction of mosquito abundance can provide significant information to authorities to design vector surveillance and prevention strategies for outbreaks. By using available data, machine learning can train and test the data, use the model results to predict future outcomes, and improve the prediction from time to time. Figure 2 shows an overview of the proposed framework.




Figure 2: Overview of the proposed framework

### 3 CONCLUSION

Due to the geographical expansion of dengue in recent years, the information gathered through surveillance is essential for risk evaluation and epidemic management, making it a vital component of dengue prevention. Surveillance activities should ideally involve ecological monitoring and social risk issues in addition to the early detection of human infections and vector control. In Malaysia, rigorous dengue control measures are implemented in certain localities based on the notification of lab-confirmed human cases and the location of dengue hotspots. Currently, house-to-house larval inspections by the health inspector, removing mosquito breeding sites, larvicide activities, and fogging are among the control measures taken by the Ministry of Health, Malaysia. The feasibility of using drone technology in mosquito breeding control programmes should be contemplated as an effective alternative tool, as it could substitute for the time-consuming conventional process of observing larval habitats. This proposed framework, which integrates drones and technologies, community assessment, and collaboration with government agencies, will add a new dimension to dengue prevention and control programmes in Malaysia and other endemic regions.


### ACKNOWLEDGEMENT

This research is supported by the Malaysian Ministry of Higher Education and Universiti Teknologi MARA (UiTM) under the Academic Training Scheme for Bumiputera (SLAB).

### REFERENCES

Ali, A., Nisar, S., Khan, M. A., Agha, S., Mohsan, H., & Noor, F. (2022). *A Privacy-Preserved Internet-of-Medical-Things Scheme for Eradication and Control of Dengue Using UAV*. *Micromachines*, 13(10), p.1702.

Amarasinghe, A., Suduwella, C., Niroshan, L., Elvitigala, C., De Zoysa, K., & Keppetiyagama, C. (2017). Suppressing dengue via a drone system. *17th International Conference on Advances in ICT for Emerging Regions, ICTer 2017 - Proceedings, 2018-Janua*, pp.202–208. <https://doi.org/10.1109/ICTER.2017.8257797>



- Annan, E., Guo, J., Angulo-Molina, A., Yaacob, W. F. W., Aghamohammadi, N., C. Guetterman, T., Yavaşoglu, S. İ., Bardosh, K., Dom, N. C., Zhao, B., Lopez-Lemus, U. A., Khan, L., Nguyen, U. S. D. T., & Haque, U. (2022). Community acceptability of dengue fever surveillance using unmanned aerial vehicles: A cross-sectional study in Malaysia, Mexico, and Turkey. *Travel Medicine and Infectious Disease*, 49, p.102360. <https://doi.org/10.1016/j.tmaid.2022.102360>
- Bravo, D. T., Lima, G. A., Alves, W. A. L., Colombo, V. P., Djogbénu, L., Pamboukian, S. V. D., Quaresma, C. C., & Araujo, S. A. de. (2021). Automatic detection of potential mosquito breeding sites from aerial images acquired by unmanned aerial vehicles. *Computers, Environment and Urban Systems*, 90, p.101692. <https://doi.org/10.1016/j.compenvurbsys.2021.101692>
- Budiharto, W., Irwansyah, E., Suroso, J. S., Chowanda, A., Ngarianto, H., & Gunawan, A. A. S. (2021). Mapping and 3D modelling using quadrotor drone and GIS software. *Journal of Big Data*, 8, pp.1-12. <https://doi.org/10.1186/s40537-021-00436-8>
- Carrasco-Escobar, G., Manrique, E., Ruiz-Cabrejos, J., Saavedra, M., Alava, F., Bickersmith, S., Prussing, C., Vinetz, J. M., Conn, J. E., Moreno, M., & Gamboa, D. (2019). High-accuracy detection of malaria vector larval habitats using drone-based multispectral imagery. *PLoS Neglected Tropical Diseases*, 13(1), pp.1–24. <https://doi.org/10.1371/journal.pntd.0007105>
- Carrasco-Escobar, G., Moreno, M., Fornace, K., Herrera-Varela, M., Manrique, E., & Conn, J. E. (2022). The use of drones for mosquito surveillance and control. *Parasites and Vectors*, 15(1), pp.1–22. <https://doi.org/10.1186/s13071-022-05580-5>
- Carrillo-Larco, R. M., Moscoso-Porras, M., Taype-Rondan, A., Ruiz-Alejos, A., & Bernabe-Ortiz, A. (2018). The use of unmanned aerial vehicles for health purposes: a systematic review of experimental studies. *Global Health, Epidemiology and Genomics*, 3, p.e13. <https://doi.org/10.1017/gheg.2018.11>
- Charmaz, K. (2014). Grounded theory in global perspective: Reviews by international researchers. *Qualitative inquiry*, 20(9), pp.1074-1084.
- Chiroli, D. M. de G., Montilha, M. M., & Samed, M. M. A. (2017). Feasibility analysis of uav in dengue control. *Independent Journal of Management & Production*, 8(2), p.319. <https://doi.org/10.14807/ijmp.v8i2.539>
- Creswell, J. W. & Poth, C. N. (2018). *Qualitative Inquiry & Research Design : Choosing Among Five Approaches*. Fourth Edition. SAGE Publications.
- Dias, T. M., Alves, V. C., Alves Junior, H. M., Pinheiro, L. F., Pontes, R. S. G., Araujo, G. M., Lima, A. A., & Prego, T. M. (2018). Autonomous detection of mosquito-breeding habitats using an unmanned aerial vehicle. *Proceedings - 15th Latin American Robotics Symposium, 6th Brazilian Robotics Symposium and 9th Workshop on Robotics in Education, LARS/SBR/WRE 2018*, pp.357–362. <https://doi.org/10.1109/LARS/SBR/WRE.2018.00070>
- Duque, E., Fonseca, G., Vieira, H., Gontijo, G., & Ishitani, L. (2019). A systematic literature review on user centered design and participatory design with older people. *IHC 2019 - Proceedings of the 18th Brazilian Symposium on Human Factors in Computing Systems*. pp.1-11. <https://doi.org/10.1145/3357155.3358471>
- Faraji, A., Haas-Stapleton, E., Sorensen, B., Scholl, M., Goodman, G., Buettner, J., Schon, S., Lefkowitz, N., Lewis, C., Fritz, B., Hoffman, C., & Williams, G. (2021). Toys or Tools? Utilization of Unmanned Aerial Systems in Mosquito and Vector Control Programs. *Journal of Economic Entomology*, 114(5), pp.1896–1909. <https://doi.org/10.1093/jee/toab107>
- Fornace, K. M., Drakeley, C. J., William, T., Espino, F., & Cox, J. (2014). Mapping infectious disease landscapes: Unmanned aerial vehicles and epidemiology. *Trends in Parasitology*, 30(11), pp.514–519. <https://doi.org/10.1016/j.pt.2014.09.001>
- Haas-Stapleton, E. J., Barretto, M. C., Castillo, E. B., Clausnitzer, R. J., & Ferdan, R. L. (2019). Assessing mosquito breeding sites and abundance using an unmanned aircraft. *Journal of the American Mosquito Control Association*, 35(3), pp.228–232. <https://doi.org/10.2987/19-6835.1>
- Hardy, A., Makame, M., Cross, D., Majambere, S., & Msellem, M. (2017). Using low-cost drones to map malaria vector habitats. *Parasites and Vectors*, 10(1), pp.1–13. <https://doi.org/10.1186/s13071-017-1973-3>
- Hardy, A., Oakes, G., Hassan, J., & Yussuf, Y. (2022). Improved Use of Drone Imagery for Malaria Vector Control through Technology-Assisted Digitizing (TAD). *Remote Sensing*, 14(2), pp.1–18. <https://doi.org/10.3390/rs14020317>
- Hasnan, S. N. A., Dom, N. C., Latif, Z. A., & Madzlan, F. (2017). Surveillance of *Aedes* mosquitoes in different residential types in central zone of Shah Alam, Selangor. *Asian Pacific Journal of Tropical Disease*, 7(6), pp.332–340. <https://doi.org/10.12980/apjtd.7.2017D7-44>
- Hiebert, B., Nouvet, E., Jeyabalan, V., & Donelle, L. (2020). The application of drones in healthcare and health-related services in North America: A scoping review. *Drones*, 4(3), pp.1–17. <https://doi.org/10.3390/drones4030030>
- Horstick, O., Runge-Ranzinger, S., Nathan, M. B., & Kroeger, A. (2010). Dengue vector-control services: How do they work? A systematic literature review and country case studies. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 104(6), pp.379–386. <https://doi.org/10.1016/j.trstmh.2009.07.027>
- Joshi, A., & Miller, C. (2021). Review of machine learning techniques for mosquito control in urban environments. *Ecological Informatics*, 61, p.101241. <https://doi.org/10.1016/j.ecoinf.2021.101241>
- Kleinman, A. (2010). Four social theories for global health. *Lancet*, 375(9725), pp.1518–1519. [https://doi.org/10.1016/S0140-6736\(10\)60646-0](https://doi.org/10.1016/S0140-6736(10)60646-0)
- Madzlan, F., Dom, N. C., Zakaria, N., Hasnan, S. N. A., Tiong, C. S., & Camalxaman, S. N. (2017). Profiling of dengue vectors breeding habitat at urban residential areas in shah alam, Malaysia. *Serangga*, 22(2), pp.185–216.
- Mehra, M., Bagri, A., Jiang, X., & Ortiz, J. (2016). Image Analysis for Identifying Mosquito Breeding Grounds. *2016 IEEE International Conference on Sensing, Communication and Networking, SECON Workshops 2016*, pp.1-6. <https://doi.org/10.1109/SECONW.2016.7746808>
- Mohd Daud, S. M. S., Mohd Yusof, M. Y. P., Heo, C. C., Khoo, L. S., Chainchel Singh, M. K., Mahmood, M. S., & Nawawi, H. (2022). Applications of drone in disaster management: A scoping review. *Science and Justice*, 62(1), pp.30–42. <https://doi.org/10.1016/j.scijus.2021.11.002>
- Mukabana, W. R., Welter, G., Ohr, P., Tingitana, L., Makame, M. H., Ali, A. S., & Knols, B. G. J. (2022). Drones for Area-Wide Larval Source Management of Malaria Mosquitoes. *Drones*, 6(7), p.180. <https://doi.org/10.3390/drones6070180>

- Muñiz-sánchez, V., Valdez-delgado, K. M., & Hernandez-lopez, F. J. (2022). Use of Unmanned Aerial Vehicles for Building a House Risk Index of Mosquito-Borne Viral Diseases. *Machines*, 10(12), p.1161. <https://doi.org/10.3390/machines10121161>
- Passos, W. L., Araujo, G. M., de Lima, A. A., Netto, S. L., & da Silva, E. A. B. (2022). Automatic detection of *Aedes aegypti* breeding grounds based on deep networks with spatio-temporal consistency. *Computers, Environment and Urban Systems*, 93, p.101754. <https://doi.org/10.1016/j.compenvurbsys.2021.101754>
- Poljak, M., & Šterbenc, A. (2020). Use of drones in clinical microbiology and infectious diseases: current status, challenges and barriers. *Clinical Microbiology and Infection*, 26(4), pp.425–430. <https://doi.org/10.1016/j.cmi.2019.09.014>
- Rahman, M. S., Pientong, C., Zafar, S., Ekalaksananan, T., Paul, R. E., Haque, U., Rocklöv, J., & Overgaard, H. J. (2021). Mapping the spatial distribution of the dengue vector *Aedes aegypti* and predicting its abundance in northeastern Thailand using machine-learning approach. *One Health*, 13, p.100358. <https://doi.org/10.1016/j.onehlt.2021.100358>
- Schenkel, J., Taelle, P., Goldberg, D., Horney, J., & Hammond, T. (2020). Identifying potential mosquito breeding grounds: Assessing the efficiency of uav technology in public health. *Robotics*, 9(4), p.91. <https://doi.org/10.3390/robotics9040091>
- Stanton, M. C., Kalonde, P., Zembere, K., Hoek Spaans, R., & Jones, C. M. (2021). The application of drones for mosquito larval habitat identification in rural environments: a practical approach for malaria control? *Malaria Journal*, 20(1), pp.1–17. <https://doi.org/10.1186/s12936-021-03759-2>
- Valdez-Delgado, K. M., Moo-Llanes, D. A., Danis-Lozano, R., Cisneros-Vázquez, L. A., Flores-Suarez, A. E., Ponce-García, G., Medina-De la Garza, C. E., Díaz-González, E. E., & Fernández-Salas, I. (2021). Field effectiveness of drones to identify potential *aedes aegypti* breeding sites in household environments from tapachula, a dengue-endemic city in southern mexico. *Insects*, 12(8), p.663. <https://doi.org/10.3390/insects12080663>
- Wyngaard, J., Rund, S. S. C., Madey, G. R., & Cleland-huang, J. (2018). for Mosquito-borne Disease Research and Control. *2018 IEEE/ACM 40th International Conference on Software Engineering: Companion (ICSE-Companion)*, pp.226–227.
- Zhang, J. (2005). Human-centered computing in health information systems Part 1: Analysis and design. *Journal of Biomedical Informatics*, 38(1), pp.1–3. <https://doi.org/10.1016/j.jbi.2004.12.002>